# Native and Non-native Plant Seedlings of Northern Arizona



Erin R. Thurston

**Ecological Restoration Institute** 

# Contents

Introduction, methods	2
About the author, acknowledgements,	
contact information	3
Grasses	_
Poaceae	4
Forbs	
Asteraceae	10
Fabaceae	11
Rosaceae	13
Scrophulariaceae	13
Shrubs	
Anacardiaceae	15
Ephedraceae	15
Rhamnaceae	16
Glossary	17
Appendix A: Species planted but not included	
due to growing difficutly	18
Appendix B: Species recommended	
but not included	18

## Introduction

Southwestern ponderosa pine forests support a wide diversity of understory vegetation, including grasses, forbs, shrubs, and cacti. Plant identification is important in monitoring following a disturbance event or in conducting ecological restoration. The purpose of this photographic guide is to assist ecologists and land managers in seedling identification and restoration monitoring.

## Methods

I performed surveys with botanists employed with the Ecological Restoration Institute (ERI) in order to determine the range of species common to the greater Flagstaff area to include in the field guide. I then obtained seeds for the selected species from botanists employed with the ERI and from seed outlets in the Flagstaff area. I performed the necessary pre-germination treatments for each species, then germinated the seeds by placing them on wet filter paper in petri dishes. I also germinated seeds by planting them in potting soil in seed flats. I manually watered the petri dishes and kept the seed flats in a Northern Arizona University greenhouse where they were watered by a timed sprinkler system. When the seedlings began to grow, I photographically documented each species, focusing on distinctive characteristics. I photographed the species with a Leica microscope fitted with an Optronics MagnaFire Sp camera, Model S99805, located in the ERI ecology lab. I also measured characteristics of each species, such as stem size and leaf size, and pressed samples of each stage.

The guide is organized in three sections--grasses, forbs, and shrubs. Each section is organized alphabetically by species. Two photographs of each plant species accompany descriptions of the species at different developmental stages (Note: certain species only have one photograph due to complications in the growing process).

This is the 1st edition of this guide, and will hopefully serve as a template for further editions. 2

## Author

Erin R. Thurston is an Undergraduate Research Assistant with the Ecological Restoration Institute (ERI) and is pursuing a Bachelor of Science degree in Environmental Science at Northern Arizona University. The creation of this identification guide was part of her undergraduate research work with the ERI.

### Acknowledgements

This project was funded through a Hooper Undergraduate Research Award and the Ecological Restoration Institute of Northern Arizona University. Special thanks to Robin Long, Luke Brandy, Steve Buckley, Mike Stoddard, and Chris McGlone.

Contact Ecological Restoration Institute PO Box 15017 Flagstaff, AZ 86011 (928) 523-7182 www.eri.nau.edu



Copyright © 2005 Erin R. Thurston, Ecological Restoration Institute 3

Aristida purpurea

Purple threeawn

**Characteristics:** Blades: narrow, flat blades, 5-7 cm tall @ 1st true leaf, 1 mm wide **Culm**: 8 mm between glume and 1st true leaf **Other**: awns on glume



Blepharoneuron tricholepis

Pine dropseed

**Characteristics:** Blades: curled tips; 1.5 cm @ 1st true leaf, 1 mm wide; 2-3 cm tall @ 2nd true leaf, 1 mm wide **Culm:** 3 mm between glume and 1st & 2nd true leaf



### Bouteloua curtipendula

Sideoats grama

**Characteristics: Blades:** curled tips, flat blades; 4.5 cm tall @ 1st true leaf, 1 mm wide, ciliate margin **Culm:** 1 cm between glume and 1st true leaf



cotyledon



1st true leaf

Bouteloua gracilis

#### Blue grama

**Characteristics: Blades:** curled tips, flat blades; 3 cm tall @ 1st true leaf, 1 mm wide **Culm:** 1 cm between glume and 1st true leaf



# Poaceae

#### Bromus tectorum

#### Cheatgrass

Characteristics: Blades: "corkscrews," but more pubescent than Elymus--over whole blade; 4.5 cm tall @ 1st true leaf, 1 mm wide Culm: 1 cm between glume and 1st true leaf Other: awn on glume; late summer/ early fall germination



#### Elymus elymoides

Squirreltail

Characteristics: Blades: "corkscrews," may be pubescent along bottom midvein, flat blade; 6-15 cm tall @ 1st true leaf, 1.5 mm wide; prominent veins Culm: reddish, 1-2 cm between glume and 1st true leaf



cotyledon



#### Festuca arizonica

Arizona fescue

Characteristics: Blades: rolled, "hair-like" blade; narrow, straight up & down; 5.5-8.5 cm tall @ 1st true leaf, 0.5 mm wide Culm: 1.2 cm between glume and 1st true leaf



cotyledon

2nd true leaf

Koeleria macrantha

Prairie Junegrass

Characteristics: Blades: 2.5 cm tall @ 1st true leaf, 0.5 mm wide; 3 cm tall @ 2nd true leaf, <1 mm wide; narrow, flat blades; short compared to Elymus & more blue-green Culm: 3 mm between glume and 1st true leaf





2nd true leaf

### Muhlenbergia montana

Mountain muhly

**Characteristics: Blades:** 2.5-3 cm tall @ 1st true leaf, 1 mm wide **Culm:** 5 mm between glume and 1st true leaf **Other:** look for prominent ligule by 1st true leaf



cotyledon w/ligule



1st true leaf

#### Poa fendleriana

#### **Muttongrass**

**Characteristics: Blades:** curled tips; 4 cm tall @ 1st true leaf, 0.5 mm wide; 6 cm tall @ 2nd true leaf, 2 mm wide **Culm:** 0.5 cm between glume and 1st true leaf **Other:** look for "RR tracks" (2 median lines) by 2nd true leaf





2nd true leaf

Sporobolus cryptandrus

Sand dropseed

**Characteristics: Blades:** "strap-shaped" leaves; 1 cm @ 1st true leaf, 1 mm wide **Culm:** 2 mm between glume and 1st true leaf





cotyledon

1st true leaf

### Centaurea diffusa

Diffuse knapweed

Characteristics: Cotyledons: petioled; 4-7 mm long, 3-4 mm wide True leaves: opposite simple leaves, "strap-shaped"; toothed margins; pubescent; 1 cm long, 4 mm wide Stem: 5-10 mm @ cotyledons; 10 mm @ 1st true leaf



cotyledons



1st true leaf

Packera multilobata

Lobeleaf groundsel

Characteristics: Cotyledons: petioled; round; 5 mm long. 4 mm wide True leaves: single, simple, alternately toothed; 2 mm long, 1.5 mm wide Stem: variable; 3-15 mm @ cotyledons; 3-15 mm @ 1st true leaves **Other:** may have purple on bottom of leaves





1st true leaf

# Asteraceae

#### <u>Senecio eremophilus</u>

Desert ragwort

**Characteristics:** Cotyledons: petioled; 5 mm long, 4 mm wide **True leaves:** single, simple; 8 mm long, 6 mm wide; alternately toothed; may be purple underneath; petiole ciliate **Stem:** may be purple; 10 mm @ cotyledon & 1st true leaves



cotyledons



2nd true leaf

# Fabaceae

### <u>Lupinus argenteus</u>

Silver lupine

**Characteristics: Cotyledons:** fleshy, round, w/distinct petiole; 5 mm long, 9 mm wide **True leaves:** single palmately compound leaf w/5 oblanceolate leaflets, each 2-4 mm long, 1 mm wide; pubescent **Stem:** 1 cm @ cotyledons; 1-3 cm @ 1st true leaves



### Oxytropis lambertii

Lambert's locoweed

**Characteristics:** Cotyledons: sessile, fleshy, oval; 8 mm long, 3-5 mm wide **True leaves:** many, linear, ciliate; 5 mm long, 1-2 mm wide **Stem:** 1-1.5 cm @ cotyledons; 1-2 cm @ 1st true leaves





true leaves

Vicia americana

American vetch

**Characteristics: Cotyledons:** sessile; 2 mm long, 1 mm wide **True leaves:** single compound leaf, split all the way to midvein; petioled, linear; 3-5 mm long, 1-2 mm wide **Stem:** 10 mm @ cotyledon; 10-15 mm @ 1st true leaves





1st true leaf

# Rosaceae

### <u>Potentilla crinita</u>

Bearded cinquefoil

**Characteristics:** Cotyledons: round, petioled; 2 mm long, 2 mm wide **True leaves:** single simple leaf; three lobes; ciliate; 2-3 mm long **Stem:** irregular stem; 3-10 mm @ cotyledon; 10 mm @ 1st true leaf



cotyledons



1st true leaf

# Scrophulariaceae

### <u>Linaria dalmatica</u>

Dalmatian toadflax

**Characteristics: Cotyledons:** sessile, pointed tip; 2 mm long, 1 mm wide **Stem:** can be purple; 10-20 mm @ cotyledon



# Scrophulariaceae

Verbascum thapsus

Common mullein

**Characteristics: Cotyledons:** sessile, ciliate, rounded tip; 1 mm long, 1 mm wide **Stem:** 5 mm @ cotyledons



cotyledons

# Shrubs- Anacardiaceae

#### <u>Rhus trilobata</u>

Skunkbush sumac

**Characteristics: Cotyledons:** sessile to subsessile, oval; 7 mm long, 4 mm wide; pink tint underneath; petioles < 1 mm long **True leaves:** single, compound, leaflets ternate w/3 lobes per leaflet, 1.5-2 cm long, 1 cm wide **Stem:** 1-1.5 cm @ cotyledons, 1.5-2.5 cm @ 1st true leaf; pink tint



cotyledons



1st true leaf

# Ephedraceae

<u>Ephedra viridis</u>

Mormon tea

**Characteristics:** Cotyledons: none (gymnosperm); 2 leaves initially emerge from seed, much like cotyledons **True leaves:** resemble pine needles; round; 1-2 cm long, 1-2 mm in diameter **Stem:** 5 mm



true leaves w/seed



true leaves

# Rhamnaceae

## Ceanothus fendleri

Fendler's ceanothus

**Characteristics: Cotyledons:** fleshy, round, sessile; 5 mm long, 3 mm wide **Stem:** 9-25 mm @ cotyledons **Other:** ciliate roots



cotyledons

### **Glossary**

Awn- A needle-like extension of the grass floret.

**Bract**- A modified leaf at the base of a flower or flower cluster. It may resemble a normal leaf, be reduced and scale-like in appearance, or large and brilliantly colored.

Ciliate- Fringed with hairs on the edge.

**Cotyledon**- The first leaf, pair or whorl of leaves to show after germinating; a seed leaf.

**Culm-** The jointed stalk or stem of grasses, usually hollow except at the nodes.

**Floret-** A small flower, usually one of a dense cluster. **Glume-** The bracteal covering of the flowers or seeds of grain and grasses.

**Ligule-** The projecting, usually tongue-like, membranous end of the lining of the leaf sheath, seen at the base of the leaf, between it and the stalk.

**Node-** Points on a stem from which leaves, shoots, or flowers grow.

Petiole- The stalk of a leaf.

Pubescent- Fuzzy, covered with down or hairs.

**Sessile-** Leaves and other plant parts that have no stalk, but instead grow directly from the stem of the plant.

**Simple leaf**- This is a leaf that consists of a single blade, unlike a compound leaf, which has two or more.

**Ternate**- Arranged in three's; especially said of a leaf with three leaflets, or of leaves whorled in three's.

Appendix A: Species planted but not included due to growing difficulty

Family	Genus	Species
Asteraceae Chenopodiaceae Poaceae Rosaceae Rosaceae	Solidago Atriplex Achnatherum Amelanchier Rosa	velutina canescens hymenoides utahensis woodsii

Appendix B: Species recommended but not included

Family	Genus	Species
Apiaceae	Lomatium	foeniculaceum
Apiaceae	Pseudocymopterus	montanus
Asteraceae	Achillea	millefolium
Asteraceae	Agoseris	glauca
Asteraceae	Antennaria	parvifolia
Asteraceae	Artemisia	carruthii
Asteraceae	Artemisia	dracunculus
Asteraceae	Artemisia	ludoviciana
Asteraceae	Bahia	dissecta
Asteraceae	Brickellia	grandiflora
Asteraceae	Cirsium	vulgare
Asteraceae	Cirsium	wheeleri
Asteraceae	Conyza	canadensis
Asteraceae	Coreopsis	tinctoria
Asteraceae	Ericameria	nauseosa
Asteraceae	Erigeron	colomexicanus
Asteraceae	Erigeron	divergens
Asteraceae	Erigeron	flagellaris
Asteraceae	Erigeron	formosissimus
Asteraceae	Erigeron	neomexicanus
	-	1

#### Family **Species** Genus speciosus Erigeron Asteraceae exilifolium Gnaphalium Asteraceae Asteraceae Helianthella quinquenervis Asteraceae Heliomeris multiflora Heterotheca villosa Asteraceae Hieracium fendleri Asteraceae bigelovii Hymenoxys Asteraceae richardsonii Hymenoxys Asteraceae Lactuca serriola Asteraceae schiedeana Laennecia Asteraceae Machaeranthera Asteraceae canescens Pseudognaphalium macounii Asteraceae Pyrrocoma Asteraceae crocea Senecio actinella Asteraceae Senecio wootonii Asteraceae Symphyotrichum falcatum Asteraceae officinale Asteraceae Taraxacum Asteraceae Tragopogon dubius Lappula Boraginaceae occidentalis Lithospermum multiflorum Boraginaceae Brassicaceae Arabis fendleri Brassicaceae Thlaspi montanum Caprifoliaceae **Symphoricarpos** oreophilus Arenaria fendleri Caryophyllaceae Caryophyllaceae Arenaria lanuginosa ssp. saxosa Chenopodiaceae Chenopodium album Chenopodiaceae Chenopodium graveolens Chenopodiaceae Chenopodium leptophyllum Convolvulaceae Convolvulus arvensis Astragalus Fabaceae castaneiformis Fabaceae Astragalus humistratus Fabaceae lanszwertii Lathyrus var. leucanthus

Fabaceae

Lupinus

19

kingii

Family	Genus	Species
Fabaceae	Melilotus	officinalis
Fabaceae	Phaseolus	angustissimus
Fabaceae	Thermopsis	montana
	Ĩ	var. montana
Fumariaceae	Corydalis	aurea
Geraniaceae	Geranium	caespitosum
Lamiaceae	Marrubium	vulgare
Linaceae	Linum	lewisii
Malvaceae	Malva	neglecta
Malvaceae	Sphaeralcea	parviflora
Nyctaginaceae	Mirabilis	decipiens
Nyctaginaceae	Mirabilis	linearis
Onagraceae	Epilobium	brachycarpum
Onagraceae	Gayophytum	diffusum
Onagraceae	Oenothera	caespitosa
Poaceae	Bromus	ciliatus
Poaceae	Bromus	inermis
Poaceae	Elymus	trachycaulus
		ssp. trachycaulus
Poaceae	Hordeum	jubatum
Poaceae	Muhlenbergia	rigens
Poaceae	Muhlenbergia	wrightii
Poaceae	Panicum	bulbosum
Poaceae	Piptochaetium	pringlei
Poaceae	Poa	pratensis
Poaceae	Schizachyrium	scoparium
Poaceae	Sporobolus	interruptus
Polemoniaceae	Ipomopsis	aggregata
Polemoniaceae	Phlox	gracilis
Polygonaceae	Eriogonum	alatum
Polygonaceae	Eriogonum	cernuum
Polygonaceae	Eriogonum	racemosum
Polygonaceae	Polygonum	douglasii
Polygonaceae	Rumex	crispus
Portulacaceae	Portulaca	oleracea 20

### Family

#### Genus

Primulaceae Ranunculaceae Rosaceae Rosaceae Rubiaceae Scrophulariaceae Scrophulariaceae Scrophulariaceae Scrophulariaceae Scrophulariaceae Scrophulariaceae Androsace Thalictrum Potentilla Potentilla Houstonia Castilleja Mimulus Orthocarpus Penstemon Penstemon Nicotiana

#### **Species**

septentrionalis fendleri hippiana subviscosa wrightii linariifolia rubellus luteus barbatus linarioides attenuata